



Lumsden, D. E., Crowe, B., Basu, A., Amin, S., Devlin, A., Dealwis, Y., ...  
Cadwgan, J. (2019). Pharmacological management of abnormal tone and  
movement in cerebral palsy. *Archives of Disease in Childhood*, 104(8), 775-  
780. <https://doi.org/10.1136/archdischild-2018-316309>

Peer reviewed version

License (if available):  
CC BY-NC

Link to published version (if available):  
[10.1136/archdischild-2018-316309](https://doi.org/10.1136/archdischild-2018-316309)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via BMJ at <http://dx.doi.org/10.1136/archdischild-2018-316309> . Please refer to any applicable terms of use of the publisher.

## **University of Bristol - Explore Bristol Research**

### **General rights**

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/pure/about/ebr-terms>

# Archives of Disease in Childhood

## Pharmacological management of abnormal tone and movement in Cerebral Palsy

Journal:	<i>Archives of Disease in Childhood</i>
Manuscript ID	archdischild-2018-316309.R2
Article Type:	Original article
Edition:	not in use
Date Submitted by the Author:	03-Mar-2019
Complete List of Authors:	Lumsden, Daniel; Guy's and St Thomas' NHS Foundation Trust, Paediatric Neurosciences Crowe, Belinda; Great Ormond Street Hospital for Children, The Neurodisability Service Basu, Anna; Royal Victoria Infirmary, Paediatric Neurology Amin, Sam; Paediatric Neurology, University Hospitals Bristol Devlin, Anita; Great North Children's Hospital, Department of Paediatric Neurology DeAlwis, Yasmin; Royal Victoria Infirmary, Paediatric Neurology Kumar, Ram; Alder Hey Children's NHS Foundation Trust, Paediatric Neurology Lodh, Rajib; Leeds Children's Hospital, Paediatric Neurorehabilitation Lundy, Claire; Royal Belfast Hospital for Sick Children, Paediatric Neurodisability Mordekar, Santosh; Sheffield Children's Hospital NHS Foundation Trust, Paediatric Neurology Smith, Martin; John Radcliffe Hospital, Paediatric Neurology Cadwgan, Jill; Guy's and St Thomas' NHS Foundation Trust, Paediatric Neurosciences
Keywords:	Neurodisability, Neurology, Pharmacology

SCHOLARONE™  
Manuscripts

1  
2  
3 Title:  
4

5 Pharmacological management of abnormal tone and movement in Cerebral  
6

7  
8 Palsy:  
9

10  
11  
12 Authors:  
13

14 Daniel E Lumsden\*<sup>1,2</sup>, Belinda Crowe<sup>3</sup>, Anna Basu<sup>4,5</sup>, Sam Amin<sup>6</sup>, Anita  
15

16 Devlin<sup>5,7</sup>, Yasmin DeAlwis<sup>5</sup>, Ram Kumar<sup>8</sup>, Rajib Lodh<sup>9</sup>, Claire Lundy<sup>10</sup>,  
17

18 Santosh R Mordekar<sup>11</sup> Martin Smith<sup>12</sup>, Jill Cadwgan<sup>1,4</sup>  
19  
20

21  
22  
23  
24 Affiliations:  
25

26 <sup>1</sup>Children's Neurosciences, Evelina London Children's Hospital, London, UK  
27

28 <sup>2</sup>Institute of Women and Children's Health, Faculty of Life Sciences and  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Medicine, King's College London, UK

<sup>3</sup>Department of Neurology, Great Ormond Street Hospital, London, UK

<sup>4</sup>Institute of Health and Society, Newcastle University, Newcastle, UK

<sup>5</sup>Department of Paediatric Neurology, Great North Children's Hospital, Royal  
Victoria Infirmary, Newcastle, UK

<sup>6</sup>Paediatric Neurology, University Hospitals Bristol, Bristol, UK

<sup>7</sup> Institute of Neuroscience, Newcastle University, Newcastle, UK

<sup>8</sup>Paediatric Neurology, Alder Hey Children's Hospital, Liverpool, UK

<sup>9</sup>Paediatric Neurorehabilitation, Leeds Children Hospital, Leeds, UK

<sup>10</sup>Paediatric Neurology, Royal Belfast Hospital for Sick Children, Belfast, UK

<sup>11</sup>Paediatric Neurology, Sheffield Children's Hospital, Sheffield, UK

<sup>12</sup>Paediatric Neurology, John Radcliffe Hospital, Oxford, UK

1  
2  
3 Word Count: (2721/2500)  
4

5 Tables: 2  
6

7 Figures: 2  
8  
9

10  
11  
12 References: 31/40  
13

14 \*Corresponding Author

15 Children's Neurosciences, Evelina London Children's Hospital, Guy's and St  
16

17 Thomas' NHS Foundation Trust  
18

19 Telephone 02071887188 Ext 88853  
20

21 [Daniel.lumsden@gstt.nhs.uk](mailto:Daniel.lumsden@gstt.nhs.uk)  
22  
23

24  
25  
26  
27  
28  
29  
30  
31 **Abstract (233/250)**  
32

33 **Background:** The evidence base to guide pharmacological management of  
34 tone and abnormal movements in Cerebral Palsy (CP) is limited, as is an  
35 understanding of routine clinical practice in the UK. We aimed to establish  
36 details of motor phenotype and current pharmacological management of a  
37 representative cohort across a network of UK tertiary centres.  
38  
39  
40  
41  
42  
43

44 **Methods:** Prospective multi-centre review of specialist motor disorder clinics  
45 at 8 UK centres, collecting data on clinical features and pharmacological  
46 management of children and young people (CYP) with CP over a single  
47 calendar month.  
48  
49  
50  
51  
52

53 **Results:** Data was collected from 275 CYP with CP reviewed over the  
54 calendar month of October 2017. Isolated dystonia or spasticity was  
55 infrequently seen, with a mixed picture of dystonia and spasticity +/-  
56  
57  
58  
59  
60

1  
2  
3 choreoathetosis identified in 194/275 (70.5%) of CYP. A co-morbid diagnosis  
4 of epilepsy was present in 103/275 (37.4%). The most commonly used  
5 medications for abnormal tone/movement were baclofen, trihexyphenidyl,  
6 gabapentin, diazepam and clonidine. Medication use appeared to be  
7 influenced separately by the presence of dystonia or spasticity. Botulinum  
8 toxin use was common (62.2%). A smaller proportion of children (12.4%) had  
9 undergone a previous neurosurgical procedure for tone/movement  
10 management.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

### 24 **Conclusions:**

25  
26 CYP with CP frequently present with a complex movement phenotype and co-  
27 morbid epilepsy. They have multiple therapy, medical and surgical  
28 management regimens. Future trials of therapeutic, pharmacological or  
29 surgical interventions in this population must adequately encompass this  
30 complexity in order to be translatable to clinical practice.  
31  
32  
33  
34  
35  
36  
37  
38  
39

### 40 **Introduction**

41  
42  
43  
44 Cerebral Palsy (CP) is the commonest form of chronic neurodisability  
45 encountered in paediatric practice and may be defined as a clinical syndrome  
46 of "*permanent disorders of the development of movement and posture,*  
47 *causing activity limitation, that are attributed to non-progressive disturbances*  
48 *that occurred in the developing fetal or infant brain*"(1). Elevated tone and  
49 involuntary movements are typically experienced by Children and Young  
50 People (CYP) with CP. Elevated tone, or hypertonia, caused by abnormal  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 muscle contraction may be due to spasticity, dystonia, rigidity, or a variable  
4  
5 combination of these pathological findings(2). The most widely adopted  
6  
7 classification system for CP is the Surveillance of Cerebral Palsy in Europe  
8  
9 (SCPE) classification(3), which classifies CYP on the basis of the  
10  
11 Predominant Motor Type. On this basis, 80-95% of CYP with CP are  
12  
13 considered to have “spastic” CP, and 5-17% the “Dyskinetic” form (dystonia  
14  
15 +/- choreoathetosis)(4, 5). One limitation of this system is that spasticity and  
16  
17 dystonia are treated as mutually exclusive, when in reality they are often co-  
18  
19 incident in children with motor disorders(6-8).  
20  
21  
22  
23  
24  
25

26 Elevated tone and involuntary movements typically interfere with independent  
27  
28 function and participation, the delivery of daily care and overall quality of life.  
29  
30 Consequently, pharmacological interventions are commonly prescribed,  
31  
32 though the evidence base to guide the clinician’s choice of most appropriate  
33  
34 medication is currently extremely limited. The recent American Academy of  
35  
36 Cerebral Palsy and Developmental Medicine (AACPD) guidelines on the  
37  
38 management of dystonia in CP concluded that there was insufficient evidence  
39  
40 to support any oral pharmacological agent, instead basing recommendations  
41  
42 entirely upon expert opinion(9). A number of recent reviews of the  
43  
44 management of Dyskinetic CP have highlighted the paucity of high quality  
45  
46 evidence in this patient group (9-11). The National Institute for Health and  
47  
48 Care Excellence in the UK has produced guidelines for the management of  
49  
50 spasticity in CYP(12), as has the Quality Standards Subcommittee of the  
51  
52 American Academy of Neurology and the Practice Committee of the Child  
53  
54 Neurology Society in the US(13). In both cases, a paucity of high quality  
55  
56  
57  
58  
59  
60

1  
2  
3 evidence was identified for enteral pharmacological interventions in particular.  
4  
5  
6  
7

8 Taken collectively, there is a pressing need for better quality evidence to  
9  
10 guide the pharmacological management of CYP with CP. CP is a  
11  
12 heterogeneous disorder, in terms of the nature of the underlying brain injury,  
13  
14 pattern of motor difficulties, functional impairments and medical co-morbidities  
15  
16 experienced. This heterogeneity creates challenges for the design of definitive  
17  
18 interventional studies, as does the current limited knowledge of  
19  
20 pharmacological management in current clinical practice beyond the  
21  
22 recommendation of experts(14, 15).  
23  
24  
25  
26  
27

28 In preparation for designing future trials of pharmacological management of  
29  
30 abnormal tone and movement in CYP with CP we performed a prospective  
31  
32 multi-centre review in the outpatient setting over a calendar month. We aimed  
33  
34 to determine:  
35  
36

- 37 • The clinical characteristics of CYP with CP seen during this snap shot  
38  
39 of clinical practice, focusing particularly on their pattern of motor  
40  
41 abnormalities  
42  
43
- 44 • Current pharmacological management of abnormal tone and  
45  
46 movement  
47  
48
- 49 • Diagnosis of co-morbid epilepsy and its management  
50  
51
- 52 • Determination of factors such as botulinum toxin use which could  
53  
54 confound future potential enrolment in pharmacological studies.  
55  
56  
57

## 58 **Method**

59  
60

1  
2  
3 A prospective evaluation of CYP with a clinical diagnosis of CP seen in  
4 tertiary paediatric neurology/neurodisability motor disorder clinics at 8 UK  
5 centres (Evelina London Children's Hospital, Great Ormond Street Hospital,  
6 Great North Children's Hospital, University Hospitals Bristol, Alder Hey  
7 Children's Hospital, Leeds Teaching Hospital, Royal Belfast Hospital for Sick  
8 Children, John Radcliffe Hospital and Sheffield Children's Hospital) was  
9 undertaken. Data was recorded from all cases seen over the calendar month  
10 October 2017, using a standardised data pro-forma. Data included details of  
11 gender, age, Gross Motor Classification System (GMFCS) level(16), Manual  
12 Ability Classification System (MACS) level(17), current medication use,  
13 diagnosis of co-morbid epilepsy, and whether the CYP had undergone a  
14 neurosurgical intervention aimed at reducing tone and/or abnormal  
15 movements - namely Deep Brain Stimulation (DBS), intrathecal baclofen  
16 (ITB) pump insertion, or Selective Dorsal Rhizotomy (SDR). The  
17 presence/absence of dystonia and/or spasticity was determined using the  
18 Hypertonia Assessment Tool (18), as outlined in Box 1. The presence of  
19 choreoathetosis as defined by the Taskforce for Childhood Movement  
20 disorder (19) was also recorded (Box 1). For all three motor abnormalities,  
21 clinicians were asked to record whether this was clinically significant. This  
22 was a subjective judgment, based on whether the recording clinician felt that  
23 the motor abnormality interfered sufficiently with function and participation,  
24 delivery of daily cares and/or quality of life to require an intervention aimed at  
25 its reduction. Descriptive statistics were used to summarise the data, with  
26 analysis performed on SPSS Version 24.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## **Results**

Data was collected from a total of 275 CYP with CP (162 male), median age 9 years (Range 1-18 years, 5 to 9 years, 25<sup>th</sup>-75<sup>th</sup> Centile). Clinico-demographic features are shown in Table 1. Both GMFCS (available for all 275 cases) and MACS levels (available for 258/275 cases) demonstrated peaks at Level II and Level V (Figure 1(a)). Details of motor phenotype are illustrated in Figure 1(b). Spasticity was identified in 242/275 (88.0%) CYP, judged clinically significant in 176/242 (72.7%). Dystonia was identified in 222/275 (80.7%) CYP; this was judged to be clinically significant in 172/222, 77.4%). Choreoathetosis was identified in 75/275 (27.2%) CYP (clinically significant in 26/75 (34.6%). Isolated spasticity was identified in only 47/275 (17.1%) CYP, isolated dystonia in 16/275 (5.8%) CYP, with no CYP demonstrating isolated choreoathetosis. A mixture of spasticity and dystonia was identified in 132/275 (48.0%) CYP, dystonia and choreoathetosis in 12/275 (4.4%) CYP and spasticity and choreoathetosis in 1/275 (0.4%) CYP. A combination of spasticity, dystonia and choreoathetosis was identified in 62/275 (22.5%). The remaining 5/275 (1.8%) CYP demonstrated no dystonia, spasticity or choreoathetosis.

Details of medication use are provided in Table 2. A total of 14 medications were identified across the cohort in use to reduce tone/abnormal movements. At the time of data collection, 98/275 (35.6%) CYP were receiving no medications for the management of abnormal tone/movement, 97/275 (35.3%) CYP 1 medication, 42/275 (15.3%) CYP 2 medications, 23/275

1  
2  
3 (8.4%) CYP 3 medications, 12/275 (4.4%) CYP 4 medications and 3/275  
4  
5 (1.1%) CYP 5 medications.  
6  
7  
8  
9

10 The most commonly used medications were baclofen (108/275, 39.3%),  
11 trihexyphenidyl (56/275, 20.4%), gabapentin (51/275, 18.5%), diazepam  
12 (36/275, 13.1%) and clonidine (28/275, 10.2%). Choice of medication  
13  
14 appeared to differ depending upon whether clinically significant spasticity or  
15  
16 dystonia was noted in the CYP (Figure 2). The use of baclofen did not seem  
17  
18 to be significantly altered by the presence or absence of dystonia, but  
19  
20 trihexyphenidyl, gabapentin and clonidine were all infrequently used when  
21  
22 dystonia was not present (Figure 2(a)). Conversely, the presence of clinically  
23  
24 significant spasticity seemed to increase the use of baclofen, but have  
25  
26 minimal impact on the use of trihexyphenidyl, gabapentin or clonidine (Figure  
27  
28 2(b)). Diazepam use also did not seem to differ with the presence of clinically  
29  
30 significant spasticity.  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 In addition to enteral medications for the management of abnormal  
41  
42 tone/movements, 171/275 (62.2%) CYP were receiving botulinum toxin  
43  
44 injections as part of their overall management.  
45  
46  
47  
48

49 An additional diagnosis of epilepsy had been made in 104/275 (37.4%) of  
50  
51 CYP, with 68/104 (65.4%) receiving 1 Anti-Epileptic Drug (AED) and 27/104  
52  
53 (26.0%) 2 or more AEDs. A total of 16 different AEDs were identified, with the  
54  
55 most commonly used being sodium valproate (51/275, 18.5%). In one case,  
56  
57 the ketogenic diet was being used to manage epilepsy.  
58  
59  
60

1  
2  
3  
4  
5 A total of 34/275 (12.4%) CYP had undergone a neurosurgical procedure – 3  
6 DBS, 6 SDR, and 25 ITB.  
7  
8  
9

## 10 11 12 **Discussion** 13

14 CP is a common condition, likely to be encountered frequently by  
15 professionals working across all settings in acute and community paediatrics.  
16  
17 Despite this, at present a higher level of evidence exists to support the use of  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
invasive neurosurgical interventions to manage tone, suitable for only a  
minority of CYP(9) compared to evidence for enteral medications, which are  
prescribed frequently for the majority of this patient group.

The data from this multi-centre prospective case review has demonstrated the  
complexity and variability of the motor phenotype, pharmacological  
management and associated co-morbidities of CYP with CP, illustrating the  
challenges in performing definitive trials of pharmacological intervention in this  
population. Data has been collected from tertiary care (specialist) clinics, and  
we believe is representative of practice in this setting in UK. Clinicians  
working in Paediatric tertiary care typically guide and advice the management  
of CYP by Secondary and Primary care teams (with active co-management of  
many CYP), and so our data is also likely to be applicable in these settings.

CYP with CP are commonly categorized on the basis of their predominant  
motor difficulty into “spastic”, “dyskinetic” or “ataxic” forms. This is an overly  
reductive approach, as in clinical practice a mixed picture is generally seen,

1  
2  
3 as underlined by our finding that 70.5% of CYP presented with dystonia and  
4 spasticity (+/- choreoathetosis). Isolated dystonia or spasticity was relatively  
5  
6 infrequently seen. Ataxic cerebral palsy is very rare, and potentially over-  
7  
8 reported(20). This is consistent with the findings of Rice and Colleagues, who  
9  
10 identified frequent dystonia in a cohort of 129 CYP with “spastic” CP(7). If the  
11  
12 results of future interventional studies are to be translatable into clinical  
13  
14 practice, they must be undertaken in children with a representative picture of  
15  
16 mixed tone/abnormal movements.  
17  
18  
19  
20  
21  
22  
23

24 The HAT is a validated means of defining whether spasticity and/or dystonia  
25  
26 are present in a child(18). No such tool exists for choreoathetosis, though the  
27  
28 recently developed Dyskinesia Impairment Scale does provide a tool for the  
29  
30 measurement of choreoathetosis separate from dystonia(21). The definitions  
31  
32 of both chorea and athetosis proposed by the SCPE(22) are very similar to  
33  
34 those proposed by the Taskforce on Childhood Movement Disorders(19), but  
35  
36 challenges with the practical recognition of these movements have been  
37  
38 raised(23). It is of note in our study the relatively infrequent recognition of  
39  
40 choreoathetosis compared to dystonia, with 2/7 centres documenting no CYP  
41  
42 with these abnormal movements over a calendar month. Further work is likely  
43  
44 to be required prior to future multi-centre interventional studies to ensure the  
45  
46 validity and reliability of the recognition of choreoathetosis between  
47  
48  
49  
50  
51 investigators.  
52  
53  
54  
55

56 Measuring the extent/severity of dystonia, spasticity and choreoathetosis in  
57  
58 CYP is challenging, and generally relies upon the application of clinically  
59  
60

1  
2  
3 applied scales. A recent review of available scales for dystonia and  
4 choreoathetosis in children with Dyskinetic Cerebral Palsy identified  
5 significant limitations, not least of which was that they have been designed to  
6 classify movement disorders at the level of body function and structures,  
7 rather than activity or participation(24). Similarly, the measurement of  
8 spasticity in the clinical setting is challenging (25-27). Put simply, there is no  
9 “perfect” tool to measure these different motor abnormalities directly, and  
10 measuring a reduction of these symptoms in isolation would fail to capture the  
11 true impact of an intervention for CYP and their families and carers(28).  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 A wide range of medications were identified in the management of abnormal  
27 tone and movement in this cohort, with the presence of spasticity and  
28 dystonia appearing to separately influence the choice of medication. There is  
29 very limited data available regarding current clinical practice for prescribing of  
30 medications for the management of abnormal tone and movement in CP or in  
31 other childhood conditions. In a retrospective analysis of 278 CYP in the UK  
32 with dystonia of different aetiology, the most commonly used medications  
33 were baclofen, trihexyphenidyl, levodopa and diazepam(15). In a prospective  
34 study of the management of dystonia in 57 CYP with CP at a tertiary care  
35 centre in Australia, baclofen and gabapentin were the most commonly used  
36 oral medications(14). Baclofen is a Gamma-aminobutyric acid (GABA) B  
37 receptor agonist, and data from our presented study confirms its widespread  
38 use in the management of hypertonia in childhood. Despite this widespread  
39 use, evidence for its efficacy is extremely limited. Recent systematic reviews  
40 of oral pharmacological management of dystonia in CP found no studies of  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 baclofen meeting their inclusion criteria (9, 11). Evidence to support the use of  
4 oral baclofen (in contrast to intrathecal baclofen) in the management of  
5 spasticity is currently limited, with contradictory findings reported(13).  
6  
7

8  
9  
10 Trihexyphenidyl, a centrally acting anti-cholinergic agent, has no role in the  
11 management of spasticity. A larger number of studies have examined the  
12 efficacy of trihexyphenidyl in the management of dystonia in CP, but results to  
13 date have been contradictory(9, 11). Whilst sufficient evidence was identified  
14 for the AAN guidelines to recommend the short-term use of diazepam (which  
15 potentiates the post-synaptic inhibitory effect of GABA) to reduce  
16 spasticity(13), longer term use for the management of spasticity and for  
17 reduction of dystonia is currently lacking(9, 11, 13). Gabapentin (an analogue  
18 of GABA) appears to be an increasingly used medication for tone reduction in  
19 CP, but published data is very limited. A retrospective study of gabapentin in  
20 69 children with dystonia (Level 4 evidence), including 25 with CP reported  
21 perceived benefits(29). Similarly, the use of clonidine (a centrally acting alpha-  
22 adrenergic agonist) in the management of hypertonicity in childhood appears  
23 to becoming more common, but published data supporting this use is limited  
24 to isolated case reports, and a single retrospective study of 33 children with  
25 dystonia of varied aetiology (level 4 evidence)(11, 30). Gabapentin is a widely  
26 used medication for the management of neuropathic pain. It remains to be  
27 determined to what extent its use in CYP is focused primarily on reduction of  
28 pain, as opposed to a primary intention to reduce elevated tone and posturing  
29 (though these two aims are not entirely distinct). Given the limited efficacy of  
30 baclofen and trihexyphenidyl experienced anecdotally in many CYP with high  
31 tone in CP, the growing use of gabapentin and clonidine is likely to reflect a  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 pragmatic approach by clinicians to exploring medications with established  
4 indications and safety profiles for new indications.  
5  
6  
7  
8  
9

10 Further adding to the complexity of management in these CYP was the  
11 comorbid diagnosis of epilepsy in almost 40%, the use of botulinum toxin in  
12 62.2%, and a previous neurosurgical intervention in 12.4%. Epilepsy is a  
13 common co-morbidity in CP, with data from the SCPE network from 1976-  
14 1998 reporting an overall incidence of 35% of CYP.(31). Higher rates of  
15 epilepsy were identified in CYP with dyskinetic or bilateral spastic CP. Many  
16 medications used primarily for the management of epilepsy may have a  
17 potentially beneficial effect on choreoathetoid movements (e.g.  
18 carbamazepine, levetiracetam)(11), posing further potential confounding  
19 factors for future definitive studies of interventions aimed at reducing  
20 abnormal tone and movements in this population. Additionally, the potential  
21 exists for AED medications to interact adversely with medications aimed at  
22 reducing tone and movement, in terms of drug metabolism and the risk of  
23 potentiating side effects (e.g. sedation/respiratory depression when baclofen  
24 and clobazam are used together).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 A number of limitations must be acknowledged to our presented study. Firstly,  
48 motor phenotype was reported by each individual centre, with no measure of  
49 reliability or reproducibility of this assessment between centres. This is  
50 particularly pertinent given that 2/7 centres identified no choreoathetosis in the  
51 CYP reported. The determination of whether identified motor abnormalities  
52 were clinically significant was subjective, and, as such, subject to the biases  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 of the individual clinicians. Data was collected on current medication use, but  
4  
5 no data was recorded on doses (units/kg), perceived or measured efficacy,  
6  
7 side effects, or previous medication use. No data was collected on the clinical  
8  
9 reasoning guiding medication choices for the CYP reviewed. Additionally, data  
10  
11 was not collected around therapeutic interventions, orthoses, and broader  
12  
13 non-neurological medical co-morbidities.  
14  
15  
16  
17  
18

### 19 Conclusion:

20  
21 This study highlights that CYP with CP represent a complex patient group,  
22  
23 with a typically mixed-motor phenotype, and often multi-modal methods of  
24  
25 management of abnormal tone and movements. A large number of  
26  
27 medications are in current use for the management of these motor difficulties,  
28  
29 most commonly baclofen and trihexyphenidyl. Currently, only very limited  
30  
31 evidence is available to support the efficacy of these medications,  
32  
33

34  
35 There is a pressing need for definitive studies to guide clinical practice, but  
36  
37 intelligent and pragmatic trial design is required. Trial design must also  
38  
39 accommodate co-morbid epilepsy, in addition to wider medical co-morbidities.  
40

41  
42 It must consider confounding interventions, including therapeutic and non-  
43  
44 pharmacological treatments. There must be multi-stakeholder engagement, to  
45  
46 include CYP, Parents, allied health professionals, medical and surgical  
47  
48 clinicians. A priority list is required to enable a collaborative approach which  
49  
50 will enable the clinical research community to move forward as a whole to  
51  
52 address this function, addressing outcomes across different levels of function,  
53  
54 across all domains of the ICF (some provisional suggestions for which are  
55  
56 outlined in Box 2). It is imperative that the design of future trials encompasses  
57  
58  
59  
60



1  
2  
3 these complexities to allow their outputs to directly inform clinical practice, and  
4  
5 lead to tangible improvements in quality of life for CYP with CP.  
6  
7  
8  
9

### 10 **Contributorship Statement**

11  
12 DEL conceptualised and designed the study, and wrote the initial manuscript  
13  
14 draft, BC and AB contributed to study design, and contributed to data  
15  
16 collection, SA, AD, TDA, RK, RL, CL, SM and MS contributed to data  
17  
18 collection and analysis, JC contributed to study design, data collection and  
19  
20 analysis. All authors reviewed and revised the manuscript, and approved the  
21  
22 final version as submitted.  
23  
24  
25

### 26 **What is already known on this topic:**

#### 27 **3 – 25 words each**

28  
29 Cerebral palsy is the most common diagnosis in children who present with  
30  
31 hypertonia; they present with a mixed pattern of tone and movement  
32  
33 abnormalities.  
34  
35  
36  
37  
38  
39

40 Despite widespread use of medications for tone management, there is little  
41  
42 evidence to inform best practice in pharmacological management of  
43  
44 hypertonia in this population.  
45  
46  
47  
48

49 Research design is limited by the complexity of presentation: the definition of  
50  
51 the population, confounding co-morbidities interventions and lack of robust  
52  
53 outcome measures.  
54  
55  
56  
57

### 58 **What this paper adds:**

## **2 – 25 words each**

This paper reports the described motor phenotype of CYP with CP presenting to specialist neurodisability services in the UK.

It is the first paper to report current prescribing practice for this population, and highlights the multiple medications used for tone management.

It highlights the prevalence of epilepsy as a co-morbidity, with anti-convulsant medication prescription, and confounding medical and surgical interventions which may affect tone management.

### **References:**

1. Rosenbaum P, Paneth N, Leviton A, *et al*. A report: the definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol Suppl*. 2007;**109**:8-14
2. Sanger TD, Delgado MR, Gaebler-Spira D, Hallett M, Mink JW. Classification and definition of disorders causing hypertonia in childhood. *Pediatrics*. 2003;**111**:e89-97
3. Prevalence and characteristics of children with cerebral palsy in Europe. *Dev Med Child Neurol*. 2002;**44**:633-40
4. Sellier E, Platt MJ, Andersen GL, Krageloh-Mann I, De La Cruz J, Cans C. Decreasing prevalence in cerebral palsy: a multi-site European population-based study, 1980 to 2003. *Dev Med Child Neurol*. 2016;**58**:85-92
5. Smithers-Sheedy H, McIntyre S, Gibson C, *et al*. A special supplement: findings from the Australian Cerebral Palsy Register, birth years 1993 to 2006. *Dev Med Child Neurol*. 2016;**58 Suppl 2**:5-10
6. Howard J, Soo B, Graham HK, *et al*. Cerebral palsy in Victoria: motor types, topography and gross motor function. *J Paediatr Child Health*. 2005;**41**:479-83
7. Rice J, Skuza P, Baker F, Russo R, Fehlings D. Identification and measurement of dystonia in cerebral palsy. *Dev Med Child Neurol*. 2017;**59**:1249-55
8. Lin JP, Lumsden DE, Gimeno H, Kaminska M. The impact and prognosis for dystonia in childhood including dystonic cerebral palsy: a clinical and demographic tertiary cohort study. *J Neurol Neurosurg Psychiatry*. 2014;**85**:1239-44
9. Fehlings D, Brown L, Harvey A, *et al*. Pharmacological and neurosurgical interventions for managing dystonia in cerebral palsy: a systematic review. *Dev Med Child Neurol*. 2018;**60**:356-66

10. Monbaliu E, Himmelmann K, Lin JP, *et al.* Clinical presentation and management of dyskinetic cerebral palsy. *Lancet Neurol.* 2017;**16**:741-9
11. Masson R, Pagliano E, Baranello G. Efficacy of oral pharmacological treatments in dyskinetic cerebral palsy: a systematic review. *Dev Med Child Neurol.* 2017;**59**:1237-48
12. Excellence NifHaC. Spasticity in children and young people with non-progressive brain disorders: Management of spasticity and co-existing motor disorders and their early musculoskeletal complications. London: National Institute for Health and Care Excellence; 2012.
13. Delgado MR, Hirtz D, Aisen M, *et al.* Practice parameter: pharmacologic treatment of spasticity in children and adolescents with cerebral palsy (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology.* 2010;**74**:336-43
14. Harvey A, Reddihough D, Scheinberg A, Williams K. Oral medication prescription practices of tertiary-based specialists for dystonia in children with cerebral palsy. *J Paediatr Child Health.* 2018;**54**:401-4
15. Lumsden DE, M.; Tomlin, S.; Lin, J-P. Medication Use in Childhood Dystonia. *Eur J Paediatr Neurol.* 2016 Jul;**20**(4):625-9
16. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol.* 1997;**39**:214-23
17. Eliasson AC, Kruminde-Sundholm L, Rosblad B, *et al.* The Manual Ability Classification System (MACS) for children with cerebral palsy: scale development and evidence of validity and reliability. *Dev Med Child Neurol.* 2006;**48**:549-54
18. Jethwa A, Mink J, Macarthur C, Knights S, Fehlings T, Fehlings D. Development of the Hypertonia Assessment Tool (HAT): a discriminative tool for hypertonia in children. *Dev Med Child Neurol.* 2010;**52**:e83-7
19. Sanger TD, Chen D, Fehlings DL, *et al.* Definition and classification of hyperkinetic movements in childhood. *Mov Disord.* 2010;**25**:1538-49
20. Smithers-Sheedy H, Badawi N, Blair E, *et al.* What constitutes cerebral palsy in the twenty-first century? *Dev Med Child Neurol.* 2014;**56**:323-8
21. Monbaliu E, Ortibus E, De Cat J, *et al.* The Dyskinesia Impairment Scale: a new instrument to measure dystonia and choreoathetosis in dyskinetic cerebral palsy. *Dev Med Child Neurol.* 2012;**54**:278-83
22. Christine C, Dolk H, Platt MJ, Colver A, Prasauskiene A, Krageloh-Mann I. Recommendations from the SCPE collaborative group for defining and classifying cerebral palsy. *Dev Med Child Neurol Suppl.* 2007;**109**:35-8
23. Eggink H, Kremer D, Brouwer OF, *et al.* Spasticity, dyskinesia and ataxia in cerebral palsy: Are we sure we can differentiate them? *Eur J Paediatric Neurol* 2017;**21**:703-6
24. Stewart K, Harvey A, Johnston LM. A systematic review of scales to measure dystonia and choreoathetosis in children with dyskinetic cerebral palsy. *Dev Med Child Neurol.* 2017;**59**:786-95
25. Love S, Gibson N, Smith N, Bear N, Blair E. Interobserver reliability of the Australian Spasticity Assessment Scale (ASAS). *Dev Med Child Neurol.* 2016;**58** Suppl 2:18-24

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
26. Scholtes VA, Becher JG, Beelen A, Lankhorst GJ. Clinical assessment of spasticity in children with cerebral palsy: a critical review of available instruments. *Dev Med Child Neurol*. 2006;**48**:64-73
27. Flamand VH, Masse-Alarie H, Schneider C. Psychometric evidence of spasticity measurement tools in cerebral palsy children and adolescents: a systematic review. *J Rehabil Med*. 2013;**45**:14-23
28. Lumsden DE, Gimeno H, Tustin K, Kaminska M, Lin JP. Interventional studies in childhood dystonia do not address the concerns of children and their carers. *Eur J Paediatr Neurol* 2015;**19**:327-36
29. Liow NY, Gimeno H, Lumsden DE, *et al*. Gabapentin can significantly improve dystonia severity and quality of life in children. *Eur J Paediatr Neurol* 2016;**20**:100-7
30. Sayer C, Lumsden DE, Kaminska M, Lin JP. Clonidine use in the outpatient management of severe secondary dystonia. *Eur J Paediatr Neurol* 2017 **21**(4):621-626
31. Sellier E, Uldall P, Calado E, *et al*. Epilepsy and cerebral palsy: characteristics and trends in children born in 1976-1998. *Eur J Paediatr Neurol* 2012;**16**:48-55

Tables:

GMFCS	Level	Number	% Cohort
	I	35	12.7
	II	51	18.5
	III	24	8.7
	IV	70	25.5
	V	95	34.5
MACS	I	37	13.5
	II	70	25.4
	III	32	11.6
	IV	48	17.5
	V	71	25.8
Dystonia	None	53	19.3
	Not Significant	50	18.2
	Clinically Significant	172	62.5
Spasticity	None	33	12.0
	Not Significant	66	24.0
	Clinically Significant	176	64.0
Choreoathetosis	None	200	72.7
	Not Significant	49	17.8
	Clinically Significant	26	9.5
Epilepsy	Yes	104	37.8
	No	171	62.2
Botulinum Toxin Program	Yes	171	62.2
	No	104	37.8
Deep Brain Stimulation	Yes	3	1.1
	No	272	98.9
Intrathecal Baclofen	Yes	25	9.1
	No	250	90.9
Selective Dorsal Rhizotomy	Yes	6	3.2
	No	269	97.8

Table 1: Clinical Features of CYP with CP reviewed in clinic.

Abbreviations used: "GMFCS" – Gross Motor Classification System, "MACS"

– Manual Ability Classification System.

Medications Type	Medication	Number	% Overall cohort
Tone/Movement	Baclofen	108	39.3
	Trihexyphenidyl	56	20.4
	Gabapentin	51	18.5
	Diazepam	36	13.1
	Clonidine	28	10.2
	Chloral Hydrate	11	4.0
	Levodopa	9	3.3
	Nitrazepam	2	0.7
	Clonazepam	2	0.7
	Midazolam	1	0.4
	Risperidone	1	0.4
	Nabiximol	1	0.4
	Amitriptyline	1	0.4
	Tizanidine	1	0.4
Epilepsy	Sodium Valproate	51	18.5
	Levetiracetam	23	8.4
	Carbamazepine	19	6.9
	Lamotrigine	11	4.1
	Topiramate	10	3.6
	Clobazam	3	1.1
	Nitrazepam	2	0.7
	Oxcarbazepine	2	0.7
	Lacosamide	2	0.7
	Vigabatrin	1	0.4
	Clonazepam	1	0.4
	Phenytoin	1	0.4
	Phenobarbitone	1	0.4

	Zonisamide	1	0.4
	Calcium Folate	1	0.4
	Perampanel	1	0.4

Table 2: Medication Use across the cohort.

### Figure Legends

#### Figure 1:

1(a) Distribution of Gross Motor Function Classification System (GMFCS) and Manual Ability Classification System (MACS) levels across the cohort. 1(b) Motor phenotype of CYP in the cohort, with presence of spasticity and/dystonia defined as per the Hypertonia Assessment Tool (HAT) and choreoathetosis as defined by the Taskforce on Childhood Motor disorders. Clinical significance of these features was based on judgment of the reporting clinicians.

#### Figure 2:

Frequency of individual medication use depending upon whether CYP demonstrated clinically significant (a) Dystonia, (b) Spasticity or (c) Choreoathetosis.

## Box 1

Spasticity	<p>Features denoting presence of spasticity according to Hypertonia Assessment Tool(10)</p> <ul style="list-style-type: none"> <li>• Velocity - dependent resistance to stretch</li> <li>• Presence of spastic catch</li> </ul>
Dystonia	<p>Features denoting presence of dystonia according to Hypertonia Assessment Tool (10)</p> <ul style="list-style-type: none"> <li>• Increased involuntary movements or postures of the designated limb with tactile stimulus of a distant body part</li> <li>• Increased involuntary movements or postures with purposeful movement of a distant body part</li> <li>• Increased tone with movement of a distant body part</li> </ul>
Choreoathetosis	<p>As defined by the Taskforce on Childhood Movement Disorders (19)</p> <p>Chorea is an ongoing random-appearing sequence of one or more discrete involuntary movements or movement fragments.</p> <p>Athetosis is a slow, continuous, involuntary writhing movement that prevents maintenance of a stable posture</p>

## Box 2

<p>Priorities to be addressed for development of future trial design of medications for the treatment of tone and movement in CP</p>
<ul style="list-style-type: none"> <li>• How to standardize a description of the motor profile of children with CP?</li> <li>• How to ensure the reliability of motor phenotyping across multiple centres and clinicians?</li> </ul>



- Accepting a single trial cannot include all children and young people with CP. How should this heterogenous cohort be divided into biologically and clinically meaningful studies for future trial designs (e.g by functional level, age, aetiology, features of motor phenotype)?
- What Outcome measures should be used? Measures must map across all domains of the ICF, and be meaningful to children and young people with CP, their parents/carers and clinicians alike?
- Over what time scale should studies be performed?
- What medications should be trialed, and compared to what? Would a placebo arm be acceptable to children and young people with CP, their parents/carers and clinicians alike? What medication would be considered the “standard of care”?
- Given the off license use of most medications in current clinical practice, what dosing range and schedules should be used for medications included in future pharmacological studies?
- To what extent can other aspects of care be standardized during the time children and young people with CP are enrolled in future studies (e.g. application of orthoses)?
- Should future trials focus on medication naïve children and young people? This would simplify study design, but would limit recruitment, preventing many children and young people with CP benefitting from and contributing to trial enrollment.




Figure 1

Figure 1:

1(a) Distribution of Gross Motor Function Classification System (GMFCS) and Manual Ability Classification System (MACS) levels across the cohort. 1(b) Motor phenotype of CYP in the cohort, with presence of spasticity and/dystonia defined as per the Hypertonia Assessment Tool (HAT) and choreoathetosis as defined by the Taskforce on Childhood Motor disorders. Clinical significance of these features was based on judgment of the reporting clinicians.

190x157mm (300 x 300 DPI)




Figure 2:  
 Frequency of individual medication use depending upon whether CYP demonstrated clinically significant (a) Dystonia, (b) Spasticity or (c) Choreoathetosis.

266x141mm (300 x 300 DPI)